

## **Clinical Policy: Dextromethorphan/Quinidine (Nuedexta)**

Reference Number: CP.PMN.93

Effective Date: 12.05.17

Last Review Date: 02.18

Line of Business: Commercial, Health Insurance Marketplace, Medicaid

[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

### **Description**

Dextromethorphan and quinidine (Nuedexta<sup>®</sup>) are a fixed-dose combination of dextromethorphan hydrobromide, an N-methyl-D-aspartate (NMDA) receptor antagonist and sigma-1 agonist, and quinidine sulfate, a CYP450 2D6 inhibitor.

### **FDA Approved Indication(s)**

Nuedexta is indicated for the treatment of pseudobulbar affect (PBA).

### **Policy/Criteria**

*Provider must submit documentation (which may include office chart notes and lab results) supporting that member has met all approval criteria*

It is the policy of health plans affiliated with Centene Corporation<sup>®</sup> that Nuedexta is **medically necessary** when the following criteria are met:

#### **I. Initial Approval Criteria**

##### **A. Pseudobulbar Affect** (must meet all):

1. Diagnosis of PBA secondary to multiple sclerosis or amyotrophic lateral sclerosis;
2. Prescribed by or in consultation with a neurologist;
3. Baseline Center for Neurologic Study-Lability Scale (CNS-LS) score  $\geq 13$ ;
4. Dose does not exceed 40 mg dextromethorphan and 20 mg quinidine per day (2 capsules/day).

**Approval duration: 12 weeks**

##### **B. Other diagnoses/indications**

1. Refer to the off-label use policy for the relevant line of business if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): CP.CPA.09 for commercial, HIM.PHAR.21 for health insurance marketplace, and CP.PMN.53 for Medicaid.

#### **II. Continued Therapy**

##### **A. Pseudobulbar Affect** (must meet all):

1. Currently receiving medication via Centene benefit or member has previously met initial approval criteria;
2. Member is responding positively to therapy, as evidenced by decrease in CNS-LS score by 3 points or greater from baseline;

- If request is for a dose increase, new dose does not exceed 40 mg dextromethorphan and 20 mg quinidine per day (2 capsules/day).

**Approval duration: 12 months**

**B. Other diagnoses/indications (must meet 1 or 2):**

- Currently receiving medication via Centene benefit and documentation supports positive response to therapy.  
**Approval duration: Duration of request or 12 months (whichever is less);** or
- Refer to the off-label use policy for the relevant line of business if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): CP.CPA.09 for commercial, HIM.PHAR.21 for health insurance marketplace, and CP.PMN.53 for Medicaid.

**III. Diagnoses/Indications for which coverage is NOT authorized:**

- Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policies – CP.CPA.09 for commercial, HIM.PHAR.21 for health insurance marketplace, and CP.PMN.53 for Medicaid or evidence of coverage documents.

**IV. Appendices/General Information**

*Appendix A: Abbreviation/Acronym Key*

ALS: amyotrophic lateral sclerosis

FDA: Food and Drug Administration

MS: multiple sclerosis

PBA: pseudobulbar affect

*Appendix B: Therapeutic Alternatives*

Not applicable

*Appendix C: General Information*

- Nuedexta was studied in 367 patients with PBA secondary to dementia, stroke, or traumatic brain injury. Although use of Nuedexta resulted in statistically significant improvement from baseline in CNS-LS scores, applicability of this data in clinical practice is limited as the study was open-label and not compared to placebo<sup>7</sup>.
- There is one randomized, double-blind, placebo-controlled phase 2 trial<sup>8</sup> evaluating the use of Nuedexta in 220 patients with aggression or agitation secondary to Alzheimer’s disease over 10 weeks. Nuedexta showed that the treatment difference in Neuropsychiatric Inventory (NPI) Agitation/Aggressive scores was -1.8 (95% CI, -2.8 to -0.7, p = 0.003) compared to placebo. Although this outcome was statistically significant, it did not meet the prespecified difference of 2.5 points. Also, unlike the total NPI score, use of the single NPI domain of agitation/aggression is not well validated as an endpoint. Additional long-term data is needed to confirm evidence of benefit and safety.

**V. Dosage and Administration**

Indication	Dosing Regimen	Maximum Dose
Pseudobulbar affect	1 capsule PO QD for the initial 7 days, then 1 capsule PO BID for maintenance	Dextromethorphan 40 mg/quinidine 20 mg (2 capsules) per day

**VI. Product Availability**

Capsules: dextromethorphan hydrobromide 20 mg and quinidine sulfate 10 mg

**VII. References**

1. Nuedexta Prescribing Information. Aliso Viejo, CA: Avanir Pharmaceuticals, Inc.: January 2015. Available at [www.nuedexta.com](http://www.nuedexta.com). Accessed October 27, 2016.
2. Miller RG, Jackson CE, Kasarskis EJ, et al. Practice parameter update: the care of the patient with amyotrophic lateral sclerosis: multidisciplinary care, symptom management, and cognitive/behavioral impairment (an evidence-based review). *American Academy of Neurology*. 2009;73:1227-1233.
3. Minden SL, Feinstein A, Kalb RC, et al. Evidence-based guideline: assessment and management of psychiatric disorders in individuals with MS. *American Academy of Neurology*. 2014;82:174-181.
4. Piro EP, Brooks BR, Cummings J, et al. Dextromethorphan plus ultra low-dose quinidine reduces pseudobulbar affect. *Ann Neurol*. November 2010; 68(5):693-702.
5. Brooks BR, Thisted RA, Appel SH, et al. Treatment of pseudobulbar affect in ALS with dextromethorphan/quinidine: a randomized trial. *Neurology*. October 26, 2004; 63(8):1364-1370.
6. Panitch HS, Thisted RA, Smith RA, et al. Randomized, controlled trial of dextromethorphan/quinidine for pseudobulbar affect in multiple sclerosis. *Ann Neurol*. May 2006; 59:780-787.
7. Hammond FM, Alexander DN, Cutler AJ, et al. PRISM II: an open-label study to assess effectiveness of dextromethorphan/quinidine for pseudobulbar affect in patients with dementia, stroke, or traumatic brain injury. *BMC Neurology*. 2016; 16:89. doi: 10.1186/s12883-016-0609-0.
8. Cummings JL, Lyketsos CG, Peskind ER, et al. Effect of dextromethorphan-quinidine on agitation in patients with Alzheimer disease dementia: a randomized clinical trial. *JAMA*. 2015;314(12):1242-1254.

Reviews, Revisions, and Approvals	Date	P&T Approval Date
Policy created	12.05.17	02.18

**Important Reminder**

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. “Health Plan” means a health plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan’s affiliates, as applicable.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

**Note:**

**For Medicaid members**, when state Medicaid coverage provisions conflict with the coverage provisions in this clinical policy, state Medicaid coverage provisions take precedence. Please refer to the state Medicaid manual for any coverage provisions pertaining to this clinical policy.

**For Health Insurance Marketplace members**, when applicable, this policy applies only when the prescribed agent is on your health plan approved formulary. Request for non-formulary drugs must be reviewed using the formulary exception policy.

©2017 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international

copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene Corporation. You may not alter or remove any trademark, copyright or other notice contained herein. Centene<sup>®</sup> and Centene Corporation<sup>®</sup> are registered trademarks exclusively owned by Centene Corporation.