

Clinical Policy: Sildenafil (Revatio)
Reference Number: HIM.PA.SP21
Effective Date: 05/17
Last Review Date:
Line of Business: Health Insurance Marketplace

[Coding Implications](#)
[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

Description

Sildenafil (Revatio®)* is a phosphodiesterase 5 (PDE5) inhibitor.

**Revatio and its generic, sildenafil, should not be confused with the brand formulation of sildenafil, Viagra.*

FDA approved indication

Revatio is indicated for the treatment of pulmonary arterial hypertension (PAH) (World Health Organization [WHO] Group I) in adults to improve exercise ability and delay clinical worsening.

Limitation of use: Adding sildenafil to bosentan therapy does not result in any beneficial effect on exercise capacity.

Policy/Criteria

Provider must submit documentation (including office chart notes and lab results) supporting that member has met all approval criteria

I. Initial Approval Criteria

A. Pulmonary Hypertension (must meet all):

1. Prescribed by or in consultation with a cardiologist or pulmonologist experienced in the diagnosis and treatment of pulmonary hypertension (PH);
2. Diagnosis of PH confirmed by right heart catheterization and classified as (a and b):
 - a. WHO Group 1: PAH (pulmonary arterial hypertension; Appendix B) and (i or ii):
 - i. Inadequate response or contraindication to acute vasodilator testing;
 - ii. Trial and failure of, or contraindication to, at least one calcium channel blocker;
 - b. WHO/NYHA Functional Class II, III or IV (Appendix C);
3. Prescribed dose of Revatio does not exceed 20 mg (oral formulations) or 10 mg (intravenous formulations) three times daily.

Approval duration: 6 months

B. Other diagnoses/indications

1. Refer to HIM.PHAR.21 if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized).

II. Continued Therapy

A. Pulmonary Hypertension (must meet all):

1. Currently receiving medication via Centene benefit or member has previously met all initial approval criteria;
2. Member is responding positively to therapy;

3. Prescribed dose of Revatio does not exceed 20 mg (oral formulations) or 10 mg (intravenous formulations) three times daily.

Approval duration: 12 months

B. Other diagnoses/indications (must meet 1 or 2):

1. Currently receiving medication via Centene benefit and documentation supports positive response to therapy; or
2. Refer to HIM.PHAR.21 if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized).

Approval duration: 12 months

III. Diagnoses/Indications for which coverage is NOT authorized:

- A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policy – HIM.PHAR.21 or evidence of coverage documents

IV. Appendices/General Information

Appendix A: Abbreviation/Acronym Key

CTEPH: chronic thromboembolic pulmonary hypertension

ETRA: endothelin receptor antagonist

FC: functional classification

IP receptor: prostacyclin receptor

NYHA: New York Heart Association

PAH: pulmonary arterial hypertension

PDE5: phosphodiesterase-5

PH: pulmonary hypertension

sGC: soluble guanylate cyclase

WHO: World Health Organization

Appendix B: Pulmonary Hypertension: WHO Classification

- Group 1: PAH (pulmonary arterial hypertension)
- Group 2: PH due to left heart disease
- Group 3: PH due to lung disease and/or hypoxemia
- Group 4: CTEPH (chronic thromboembolic pulmonary hypertension)
- Group 5: PH due to unclear multifactorial mechanisms

Appendix C: Pulmonary Hypertension: WHO/NYHA Functional Classes (FC)

Treatment Approach*	FC	Status at Rest	Tolerance of Physical Activity (PA)	PA Limitations	Heart Failure
Monitoring for progression of PH and treatment of co-existing conditions	I	Comfortable at rest	No limitation	Ordinary PA does not cause undue dyspnea or fatigue, chest pain, or near syncope.	

Treatment Approach*	FC	Status at Rest	Tolerance of Physical Activity (PA)	PA Limitations	Heart Failure
Advanced treatment of PH with PH-targeted therapy - see Appendix D**	II	Comfortable at rest	Slight limitation	Ordinary PA causes undue dyspnea or fatigue, chest pain, or near syncope.	
	III	Comfortable at rest	Marked limitation	Less than ordinary PA causes undue dyspnea or fatigue, chest pain, or near syncope.	
	IV	Dyspnea or fatigue may be present at rest	Inability to carry out any PA without symptoms	Discomfort is increased by any PA.	Signs of right heart failure

*PH supportive measures may include diuretics, oxygen therapy, anticoagulation, digoxin, exercise, pneumococcal vaccination. **Advanced treatment options also include calcium channel blockers.

Appendix D: Pulmonary Hypertension: Targeted Therapies

Mechanism of Action	Drug Class	Drug Subclass	Drug	Brand/Generic Formulations
Reduction of pulmonary arterial pressure through vasodilation	Prostacyclin* pathway agonist <i>*Member of the prostanoid class of fatty acid derivatives.</i>	Prostacyclin	Epoprostenol	Veletri (IV) Flolan (IV) Flolan generic (IV)
		Synthetic prostacyclin analog	Treprostinil	Orenitram (oral tablet) Remodulin (IV) Tyvasco (inhalation)
			Iloprost	Ventavis (inhalation)
		Non-prostanoid prostacyclin receptor (IP receptor) agonist	Selexipag	Uptravi (oral tablet)
	Endothelin receptor antagonist (ETRA)	Selective receptor antagonist	Ambrisentan	Letairis (oral tablet)
		Nonselective dual action receptor antagonist	Bosentan	Tracleer (oral tablet)
	Macitentan		Opsummit (oral tablet)	
	Nitric oxide-cyclic guanosine monophosphate enhancer	Phosphodiesterase type 5 (PDE5) inhibitor	Sildenafil	Revatio (IV, oral tablet, oral suspension)
			Tadalafil	Adcirca (oral tablet)
		Guanylate cyclase stimulant (sGC)	Riociguat	Adempas (oral tablet)

V. References

1. Revatio prescribing information. New York, NY: Pfizer Inc.; April 2015. Available at <http://labeling.pfizer.com/ShowLabeling.aspx?id=645>. Accessed February 10, 2017.

2. McLaughlin VV, Archer SL, Badesch DB, et al. ACCF/AHA 2009 expert consensus document on pulmonary hypertension: A report of the American College of Cardiology Foundation Task Force on Expert Consensus Documents and the American Heart Association - developed in collaboration with the American College of Chest Physicians, American Thoracic Society, Inc., and the Pulmonary Hypertension Association. *J Am Coll Cardiol*. 2009; 53(17): 1573-1619.
3. Taichman D, Ornelas J, Chung L, et. al. CHEST guideline and expert panel report: Pharmacologic therapy for pulmonary arterial hypertension in adults. *Chest*. 2014; 146(2): 449-475.
4. Abman SH, Hansmann G, Archer SL, et al. Pediatric pulmonary hypertension: Guidelines from the American Heart Association and American Thoracic Society. *Circulation*. 2015 Nov 24; 132(21): 2037-99.
5. Kim NH, Delcroix M, Jenkins DP, et al. Chronic thromboembolic pulmonary hypertension. *J Am Coll Cardiol* 2013; 62(25): Suppl D92-99.
6. Rubin LJ and Hopkins W. Overview of pulmonary hypertension in adults. In: UpToDate, Waltham, MA: Walters Kluwer Health; 2017. Available at UpToDate.com. Accessed February 14, 2017.
7. Rubin LJ and Hopkins W. Clinical features and diagnosis of pulmonary hypertension in adults. In: UpToDate, Waltham, MA: Walters Kluwer Health; 2017. Available at UpToDate.com. Accessed February 14, 2017.
8. Hopkins W and Rubin LJ. Treatment of pulmonary hypertension in adults. In: UpToDate, Waltham, MA: Walters Kluwer Health; 2017. Available at UpToDate.com. Accessed February 14, 2017.
9. Fedullo PF. Clinical manifestations and diagnosis of chronic thromboembolic pulmonary hypertension. In: UpToDate, Waltham, MA: Walters Kluwer Health; 2017. Available at UpToDate.com. Accessed February 14, 2017.
10. Fedullo PF. Overview of the treatment of chronic thromboembolic pulmonary hypertension. In: UpToDate, Waltham, MA: Walters Kluwer Health; 2017. Available at UpToDate.com. Accessed February 14, 2017.
11. Fedullo PF. Chronic thromboembolic pulmonary hypertension: Surgical treatment. In: UpToDate, Waltham, MA: Walters Kluwer Health; 2017. Available at UpToDate.com. Accessed February 14, 2017.
12. Fedullo PF. Chronic thromboembolic pulmonary hypertension: Medical treatment. In: UpToDate, Waltham, MA: Walters Kluwer Health; 2017. Available at UpToDate.com. Accessed February 14, 2017.

Reviews, Revisions, and Approvals	Date	P&T Approval Date
Policy created.	02/17	05/17

Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional

organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. “Health Plan” means a health plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan’s affiliates, as applicable.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

Note: For Medicaid members, when state Medicaid coverage provisions conflict with the coverage provisions in this clinical policy, state Medicaid coverage provisions take precedence. Please refer to the state Medicaid manual for any coverage provisions pertaining to this clinical policy.

©2017 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene Corporation. You may not alter or remove any trademark, copyright or other notice contained herein. Centene[®] and Centene Corporation[®] are registered trademarks exclusively owned by Centene Corporation.