

Clinical Policy: Temeirolimus (Torisel)

Reference Number: CP.PHAR.324

Effective Date: 03.17

Last Review Date: 11.17

Line of Business: Medicaid

[Coding Implications](#)
[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

Description

Temeirolimus for injection (Torisel[®]) is a kinase inhibitor.

FDA Approved Indication(s)

Torisel is indicated for the treatment of advanced renal cell carcinoma (RCC).

Policy/Criteria

Provider must submit documentation (which may include office chart notes and lab results) supporting that member has met all approval criteria

It is the policy of health plans affiliated with Centene Corporation[®] that Torisel is **medically necessary** when the following criteria are met:

I. Initial Approval Criteria

A. Renal Cell Carcinoma (must meet all):

1. Diagnosis of advanced RCC (i.e., relapsed, metastatic or stage IV disease; clear cell or non-clear cell histology);
2. At the time of request, member does not have bilirubin > 1.5 times the upper limit of normal;
3. Dose does not exceed 25 mg once weekly (50 mg once weekly if member is on concomitant strong CYP3A4 inducer such as dexamethasone, phenytoin, carbamazepine, rifampin, rifabutin, rifampacin, phenobarbital).

Approval duration: 6 months

B. Endometrial Carcinoma (off-label) (must meet all):

1. Diagnosis of endometrial carcinoma;
2. Use is as a single agent;
3. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (prescriber must submit supporting evidence).

Approval duration: 6 months

C. Soft Tissue Sarcoma (off-label) (must meet all):

1. Diagnosis of PEComa, recurrent angiomyolipoma, or lymphangiomyomatosis;
2. Use is as a single agent;

CLINICAL POLICY

Temsirolimus

3. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

Approval duration: 6 months

D. Other diagnoses/indications

1. Refer to CP.PHAR.57 if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized).

II. Continued Therapy

A. All Indications in Section I (must meet all):

1. Currently receiving medication via Centene benefit or member has previously met all initial approval criteria;
2. Member is responding positively to therapy (e.g., no disease progression, no significant toxicity);
3. If request is for a dose increase, request meets one of the following (a or b):
 - a. For RCC: New dose does not exceed 25 mg once weekly (*50 mg once weekly if member is on concomitant strong CYP3A4 inducer such as dexamethasone, phenytoin, carbamazepine, rifampin, rifabutin, rifampacin, phenobarbital*);
 - b. For endometrial carcinoma and soft tissue sarcoma: New dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

Approval duration: 12 months

B. Other diagnoses/indications (must meet 1 or 2):

1. Currently receiving medication via Centene benefit and documentation supports positive response to therapy.

Approval duration: Duration of request or 6 months (whichever is less); or

2. Refer to CP.PHAR.57 if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized).

III. Diagnoses/Indications for which coverage is NOT authorized:

- A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policy – CP.PHAR.57 evidence of coverage documents.

IV. Appendices/General Information

Appendix A: Abbreviation/Acronym Key

FDA: Food and Drug Administration

RCC: renal cell carcinoma

V. Dosage and Administration

Indication	Dosing Regimen	Maximum Dose
RCC	Administer 25 mg over a 30-60 minute period once a week. Treat until disease progression or unacceptable toxicity.	50 mg per week

CLINICAL POLICY

Temsirolimus

	If concomitant strong CYP3A4 inducer [e.g. dexamethasone, phenytoin, carbamazepine, rifampin, rifabutin, rifampacin, phenobarbital]): May increase to 50 mg once a week.	
--	---	--

VI. Product Availability

Kit: single-use vial 25 mg/mL temsirolimus; 1.8 mL of diluent vial

VII. References

1. Torisel Prescribing Information. Philadelphia, PA: Pfizer, Inc.; July 2017. Available at <http://labeling.pfizer.com/showlabeling.aspx?id=490>. Accessed August 30, 2017.
2. Temsirolimus. In: National Comprehensive Cancer Network Drugs and Biologics Compendium. Available at NCCN.org. Accessed August 30, 2017.

Coding Implications

Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

HCPCS Codes	Description
J9330	Injection, temsirolimus, 1 mg

Reviews, Revisions, and Approvals	Date	P&T Approval Date
Policy split from CP.PHAR.182 Excellus Oncology.	02.17	03.17
Policy converted to new template. Annual Review. Safety criteria was applied according to the safety guidance discussed at CPAC and endorsed by Centene Medical Affairs. Authorization limits extended from 3 and 6 months to 6 and 12 months for initial and continued approval, respectively. Added criteria for NCCN 2A and above recommended off-label uses: Endometrial cancer and soft tissue sarcoma.	08.30.17	11.17

Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in

CLINICAL POLICY

Temsirolimus

developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. “Health Plan” means a health plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan’s affiliates, as applicable.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

Note: For Medicaid members, when state Medicaid coverage provisions conflict with the coverage provisions in this clinical policy, state Medicaid coverage provisions take precedence. Please refer to the state Medicaid manual for any coverage provisions pertaining to this clinical policy.

CLINICAL POLICY**Temsirolimus**

©2017 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene Corporation. You may not alter or remove any trademark, copyright or other notice contained herein. Centene[®] and Centene Corporation[®] are registered trademarks exclusively owned by Centene Corporation.